


Management of Carpal Tunnel Syndrome

ANTHONY J. VIERA, LCDR, MC, USNR, Naval Hospital, Jacksonville, Florida

Carpal tunnel syndrome affects approximately 3 percent of adults in the United States. Pain and paresthesias in the distribution of the median nerve are the classic symptoms. While Tinel's sign and a positive Phalen's maneuver are classic clinical signs of the syndrome, hypalgesia and weak thumb abduction are more predictive of abnormal nerve conduction studies. Conservative treatment options include splinting the wrist in a neutral position and ultrasound therapy. Orally administered corticosteroids can be effective for short-term management (two to four weeks), but local corticosteroid injections may improve symptoms for a longer period. A recent systematic review demonstrated that nonsteroidal anti-inflammatory drugs, pyridoxine, and diuretics are no more effective than placebo in relieving the symptoms of carpal tunnel syndrome. If symptoms are refractory to conservative measures or if nerve conduction studies show severe entrapment, open or endoscopic carpal tunnel release may be necessary. Carpal tunnel syndrome should be treated conservatively in pregnant women because spontaneous postpartum resolution is common. (Am Fam Physician 2003;68:265-72, 279-80. Copyright© 2003 American Academy of Family Physicians.)

A patient information *A patient information* *handout on carpal tunnel syndrome, written by the author of this article, is provided on page 279.*



Members of various family practice departments develop articles for "Practical Therapeutics." This article is one in a series coordinated by the Department of Family Medicine at Naval Hospital, Jacksonville, Fla. Guest editor of the series is Anthony J. Viera, LCDR, MC, USNR.

See page 204 for definitions of strength-of-evidence levels.

Carpal tunnel syndrome, the most common focal peripheral neuropathy, results from compression of the median nerve at the wrist.¹ The syndrome affects an estimated 3 percent of adult Americans and is approximately three times more common in women than in men.² High prevalence rates have been reported in persons who perform certain repetitive wrist motions, but the significance of this relationship continues to be challenged. Although 30 percent of frequent computer users complain of hand paresthesias, only 10 percent meet clinical criteria for carpal tunnel syndrome, and nerve conduction studies are abnormal in only 3.5 percent of these persons.³

Family physicians frequently encounter patients who may have carpal tunnel syndrome. This article reviews the clinical features, diagnosis, and treatment of this relatively common condition.

Clinical Features

The classic symptoms of carpal tunnel syndrome are pain, numbness, and tingling in the distribution of the median

nerve (Figure 1), although numbness in all fingers may be a more common presentation.⁴ Symptoms are usually worse at night and can awaken patients from sleep. To relieve the symptoms, patients often "flick" their wrist as if shaking down a thermometer (flick sign).

In patients with carpal tunnel syndrome, pain and paresthesias may radiate to the forearm, elbow, and shoulder. Decreased grip strength may result in loss of dexterity, and thenar muscle atrophy may develop if the syndrome is severe.




FIGURE 1. Innervation of the hand (palmar and dorsal views).

ILLUSTRATIONS BY RENEE L. CANNON

Patients with carpal tunnel syndrome often have bilateral symptoms, but one hand tends to be more significantly affected.

TABLE 1
Causes and Contributing Factors in Carpal Tunnel Syndrome

Aberrant anatomy	Metabolic conditions
Anomalous flexor tendons	Acromegaly
Congenitally small carpal canal	Amyloidosis
Ganglionic cysts	Diabetes
Lipoma	Hypothyroidism or hyperthyroidism
Proximal lumbrical muscle insertion	Increased canal volume
Thrombosed artery	Congestive heart failure
Infections	Edema
Lyme disease	Obesity
Mycobacterial infection	Pregnancy
Septic arthritis	
Inflammatory conditions	
Connective tissue disease	
Gout or pseudogout	
Nonspecific flexor tenosynovitis*	
Rheumatoid arthritis	

*—Most common cause of carpal tunnel syndrome.

Adapted with permission from Von Schroeder HP, Botte MJ. Carpal tunnel syndrome. Hand Clin 1996;12:644, with additional information from Stevens JC, Beard CM, O'Fallon WM, Kurland LT. Conditions associated with carpal tunnel syndrome. Mayo Clin Proc 1992;67:541-8.

The Author

ANTHONY J. VIERA, LCDR, MC, USNR, is a staff family physician in the family practice residency program at Naval Hospital, Jacksonville, Fla., and assistant professor of family medicine at the Uniformed Services University of the Health Sciences F. Edward Hébert School of Medicine, Bethesda, Md. Dr. Viera received his medical degree from the Medical University of South Carolina College of Medicine, Charleston, and completed a family practice residency at Naval Hospital, Jacksonville.

Address correspondence to Anthony J. Viera, LCDR, MC, USNR, Family Practice Department, Naval Hospital, 2080 Child St., Jacksonville, FL 32214. Reprints are not available from the author.

Although one hand typically has more severe symptoms, both hands often are affected.

Nonspecific flexor tenosynovitis is the most common cause of carpal tunnel syndrome. However, many conditions, including aberrant anatomy, infections, inflammatory diseases, and metabolic disorders, can cause or exacerbate the syndrome (*Table 1*).^{5,6}

Diagnosis

One systematic review⁷ evaluated the effectiveness of findings from the history and physical examination in predicting positive nerve conduction studies. The most highly predictive findings were symptom location (i.e., a classic or probable pattern marked on hand symptom diagrams), hypalgesia (diminished sensitivity to pain along the palmar aspect of the index finger), and weak thumb abduction.

The principal clinical tests for carpal tunnel syndrome are Phalen's maneuver and Tinel's sign. Phalen's maneuver is positive when flexing the wrist to 90 degrees for one minute elicits symptoms in the median nerve distribution. Tinel's sign is positive when tapping over the carpal tunnel elicits symptoms in the distribution of the median nerve. Sensory findings in carpal tunnel syndrome also may be elicited by two-point discrimination, vibration, and monofilament testing.

Whether carpal tunnel syndrome is a clinical or electrophysiologic diagnosis remains somewhat controversial. In one study of 2,466 persons in a general population,² 354 (14.4 percent) reported pain, numbness, or tingling in the distribution of the median nerve. Nerve conduction studies confirmed the presence of median nerve neuropathy in approximately 45 percent of these symptomatic patients. Interestingly, nerve conduction studies were negative in almost one third of "clinically certain" patients but positive in nearly one third of "clinically uncertain" patients. Of the 125 asymptomatic patients who were examined (control group), 23 (18.4 percent) were found to have median nerve neuropathy on nerve conduction testing.

Consensus committees from the American Academy of Neurology, American Association of Electrodiagnostic Medicine, and American Academy of Physical Medicine and Rehabilitation recognize nerve conduction studies as the diagnostic standard for carpal tunnel syndrome.^{7,8}

Treatment

GENERAL MEASURES

Patients with carpal tunnel syndrome should avoid repetitive wrist and hand motions that may exacerbate symptoms or make symptom relief difficult to achieve. If possible, they should not use vibratory tools (e.g., jackhammers, floor sanders), because the motion of these tools can make their symptoms worse.⁶

Ergonomic measures to relieve symptoms depend on the motion that needs to be minimized. Patients who work on computers, for example, may benefit from improved wrist positioning or the use of wrist supports, although the latter is controversial. Wrist splints may be helpful for patients in other professions that require repetitive wrist motion.

In addition to wrist splinting, conservative treatments include oral corticosteroid therapy and local corticosteroid injections. Approximately 80 percent of patients with carpal tunnel syndrome initially respond to conservative treatment; however, symptoms recur in 80 percent of these patients after one year.⁹

One group of investigators developed an approach that uses risk factors to predict the likelihood of success for conservative treatment of carpal tunnel syndrome (Figure 2).¹⁰ Surgery should be considered when carpal tunnel syndrome does not respond to conservative measures.

WRIST SPLINTS

Splinting the wrist at a neutral angle helps to decrease repetitive flexion and rotation, thereby relieving mild soft tissue swelling or

Nerve conduction studies are the recognized diagnostic standard for carpal tunnel syndrome.

tenosynovitis. Splinting is probably most effective when it is applied within three months of the onset of symptoms.¹¹

The optimal splinting regimen depends on the patient's symptoms and preferences. Nightly splint use is recommended to prevent prolonged wrist flexion or extension.¹² When worn at night for four weeks, a specially designed wrist brace was found to be more effective than no treatment in relieving the symptoms of carpal tunnel syndrome (Figure 3).¹³ This brace has not yet been compared with traditional splints.

Predicting the Outcome of Conservative Treatment for Carpal Tunnel Syndrome

*Score 1 point for each "yes" answer and zero for each "no" answer. See the scoring key for the predicted successful outcome of conservative treatment.**

- | | | |
|---|-----------|----------|
| 1. Have symptoms been present for more than 10 months? | Yes _____ | No _____ |
| 2. Does the patient have constant paresthasias? | Yes _____ | No _____ |
| 3. Does the patient have flexor tenosynovitis ("triggering" of the digits)? | Yes _____ | No _____ |
| 4. Is Phalen's maneuver positive within less than 30 seconds? | Yes _____ | No _____ |
| 5. Is the patient older than 50 years? | Yes _____ | No _____ |

SCORING KEY: zero points = 65% success rate; 1 point = 41.4% success rate; 2 points = 16.7% success rate; 3 points = 6.8% success rate; 4 or 5 points = 0% success rate.

*—Outcome rates are based on the use of wrist splinting and nonsteroidal anti-inflammatory drugs; success rates may be higher with oral corticosteroid therapy or local corticosteroid injection.

FIGURE 2. An approach to predicting the outcome of conservative treatment in patients with carpal tunnel syndrome.

Adapted with permission from Kaplan SJ, Glickel SZ, Eaton RG. Predictive factors in the non-surgical treatment of carpal tunnel syndrome. J Hand Surg [Br] 1990;15:108.

Clinical trials in patients with carpal tunnel syndrome have shown that oral corticosteroid therapy, local injection of a corticosteroid plus an anesthetic, and ultrasound therapy provide greater symptom relief than placebo.




FIGURE 3. Manu hand brace for the conservative treatment of carpal tunnel syndrome (palmar and dorsal views). This specially designed brace provides gentle pressure to the heads of the metacarpal bones while stretching the third and fourth fingers.

Some patients choose to wear a wrist splint all of the time. Compared with nighttime-only splint use, full-time use has been shown to provide greater improvement of symptoms and electrophysiologic measures; however, compliance with full-time use is more difficult.¹⁴ [Evidence level B, uncontrolled trial]

ORAL MEDICATIONS

Diuretics, nonsteroidal anti-inflammatory drugs (NSAIDs), pyridoxine (vitamin B₆), and orally administered corticosteroids have been used with varying degrees of success in patients with carpal tunnel syndrome. Unfortunately, few high-quality trials have evaluated these treatments.

A recent systematic review¹⁵ of conservative treatments for carpal tunnel syndrome

found 14 randomized controlled trials (RCTs), including five trials of oral medications (Table 2).¹⁵⁻²⁰ The authors concluded that NSAIDs, diuretics, and pyridoxine are no more effective than placebo in relieving the symptoms of carpal tunnel syndrome.¹⁵ [Evidence level A, systematic review of RCTs] Nevertheless, some investigators²¹ recommend NSAID therapy as an adjunct to wrist splinting and ergonomic adjustments in patients with mild to moderate carpal tunnel syndrome.

Orally administered corticosteroids have been shown to be more effective than NSAIDs or diuretics in the short-term treatment of carpal tunnel syndrome. The optimal corticosteroid dosage remains to be determined. In one prospective, randomized, double-blind, placebo-controlled trial (73 patients), global symptom scores for carpal tunnel syndrome were significantly improved at two weeks and four weeks in patients randomized to receive prednisolone in a dosage of 20 mg per day for two weeks, followed by 10 mg per day for two weeks.¹⁷ [Evidence level B, lower quality RCT] No major adverse effects were noted. The study also found that NSAIDs and diuretics conferred no greater benefit than placebo.

LOCAL INJECTION

Combined injection of a corticosteroid and a local anesthetic into or proximal to the carpal tunnel can be used in patients with mild to moderate carpal tunnel syndrome. Such injections can be diagnostic as well as therapeutic. While most studies evaluating local injection have been retrospective or uncontrolled, two recent systematic reviews of RCTs concluded that local corticosteroid injection provides greater clinical improvement at one month compared with placebo.^{15,22} [References 15 and 22—Evidence level A, systematic reviews of RCTs]

A randomized, double-blind trial²³ compared oral corticosteroid therapy and local corticosteroid injection in 60 patients with

TABLE 2
Randomized Controlled Trials of Oral Medications
in the Treatment of Carpal Tunnel Syndrome

Reference	Medication(s)	Conclusions
Spooner, et al. ¹⁶	Pyridoxine (vitamin B ₆), placebo	Pyridoxine is no more effective than placebo. [Evidence level A, RCT]
Chang, et al. ¹⁷	NSAID, diuretic, corticosteroid (oral), placebo	Oral corticosteroid therapy is beneficial for short-term symptom management (4 weeks); diuretics and NSAIDs are no more effective than placebo. [Evidence level B, lower quality RCT]
Pal, et al. ¹⁸	Diuretic, placebo	Diuretic is no more effective than placebo. [Evidence level B, lower quality RCT]
Stransky, et al. ¹⁹	Pyridoxine, placebo	Pyridoxine is no more effective than placebo. [Evidence level B, lower quality RCT]
Herskovitz, et al. ²⁰	Corticosteroid (oral), placebo	Oral corticosteroid therapy is effective for short-term management (2 weeks, but not 4 weeks). [Evidence level B, lower quality RCT]

RCT = randomized controlled trial; NSAID = nonsteroidal anti-inflammatory drug.

Adapted with permission from Gerritsen AA, de Krom MC, Struijs MA, Scholten RJ, de Vet HC, Bouter LM. Conservative treatment options for carpal tunnel syndrome: a systematic review of randomised controlled trials. *J Neurol* 2002;249:276-7.

electrophysiologically confirmed carpal tunnel syndrome. In this study, 30 patients were treated with local injection of 15 mg of methylprednisolone plus oral placebo; the other 30 patients received 25 mg of orally administered prednisolone for 10 days plus a saline injection. Mean global symptom scores for the two groups did not differ significantly at two weeks. At eight and 12 weeks, however, significant improvement was evident only in the methylprednisolone injection group.

A traditional method for direct injection into the carpal tunnel is shown in *Figure 4*. Alternatively, injection can be accomplished by advancing the needle through the flexor retinaculum using a perpendicular approach.^{24,25}

Direct injection into the carpal tunnel by either method carries the potential for needle injury to the median nerve, intratendinous injection and tendon rupture, or dysesthesias (secondary to intrafascicular injection) that



FIGURE 4. Method of injecting directly into the carpal tunnel. After the hand is positioned on a rolled towel, a mixture of 10 to 20 mg of lidocaine (Xylocaine) without epinephrine and 20 to 40 mg of methylprednisolone acetate (Depo-Medrol) or similar corticosteroid preparation is injected with a 25-gauge needle at the distal wrist crease (or 1 cm proximal to it). Injection occurs along the right side of the palmaris longus tendon, which can be identified by having the patient pinch the thumb and fifth fingers together while slightly flexing the wrist. If the palmaris longus tendon cannot be identified, the needle is inserted slightly ulnar to the midline. The needle is angled downward at a 45-degree angle toward the tip of the middle finger and advanced 1 to 2 cm as it traverses the flexor retinaculum. Discomfort in the fingers should prompt repositioning of the needle.




FIGURE 5. Method of injecting proximal to the carpal tunnel. Using a 3-cm-long, 0.7-mm needle introduced at a 10- to 20-degree angle, a mixture of 10 mg of lidocaine (Xylocaine) and 40 mg of methylprednisolone is injected at the distal wrist crease between the tendons of the palmaris longus and flexor carpi radialis muscles. The mixture is introduced as a bolus and massaged toward the carpal tunnel. The needle should be advanced slowly and repositioned if resistance is encountered or the patient reports pain or paresthesias in the fingers.

may persist for months.²⁶ An alternative approach is to place the injection proximal to the carpal tunnel, rather than directly in it (*Figure 5*). This approach lowers the risk of damage to the median nerve and theoretically treats concomitant swelling at the volar side of the forearm. With any method, injection of corticosteroid into the median nerve must be avoided.

In one double-blind, placebo-controlled trial,²⁷ 60 patients older than 18 years with electrophysiologically confirmed carpal tunnel syndrome and symptoms for longer than three months were randomized to receive 10 mg of lignocaine (lidocaine) or 10 mg of lignocaine plus 40 mg of methylprednisolone. In all patients, the injection was administered proximal to the carpal tunnel. Improvement was defined as no symptoms or minor symptoms requiring no further treatment. At one month, symptoms were improved in 23 (77 percent) of the 30 patients in the corticosteroid-treated group compared with six (20 percent) of the 30 patients in the control group ($P < .001$); the number needed to treat

was 1.8.²⁷ [Evidence level A, RCT] No side effects were reported. Superior symptom relief for corticosteroid injection compared with placebo (lidocaine only) injection was not adequately demonstrated beyond one month of follow-up. All injections in the study were performed by one neurologist, which may limit the reproducibility of the results.

Splinting is generally recommended after local corticosteroid injection.^{5,25} If the first injection is successful, a repeat injection can be considered after a few months.⁹ Surgery should be considered if a patient needs more than two injections.⁹ While a poor response to local corticosteroid injection does not predict a poor surgical result, a favorable response has been associated with a satisfactory surgical outcome.⁵

ULTRASOUND THERAPY

Ultrasound therapy may be beneficial in the longer term management of carpal tunnel syndrome. A double-blind, randomized trial²⁸ found that compared with “sham ultrasound” treatment (control), 20 sessions of carpal tunnel ultrasound therapy administered over approximately seven weeks resulted in significantly greater improvement of symptoms at two weeks, seven weeks, and six months. However, a smaller study showed no benefit for this treatment.²⁹ More studies are needed to confirm the usefulness of ultrasound therapy for carpal tunnel syndrome.

Additional randomized trials comparing conservative therapies for carpal tunnel syndrome would be useful in helping physicians select appropriate treatments for individual patients.

SURGERY

Carpal tunnel release surgery should be considered in patients with symptoms that do not respond to conservative measures and in patients with severe nerve entrapment as evidenced by nerve conduction studies, thenar atrophy, or motor weakness. It is important to note that surgery may be effective

even if a patient has normal nerve conduction studies.^{30,31}

Two thirds of a community-based cohort of patients who underwent carpal tunnel release reported being completely or very satisfied with the outcome at six, 18, and 30 months after surgery.³² A recent study³³ showed that open carpal tunnel release resulted in better outcomes than wrist splinting.

Carpal tunnel release surgery is an outpatient procedure that is performed using regional anesthesia. The traditional surgical approach uses a long palmar curvilinear incision to facilitate division of the transverse carpal ligament and its overlying structures. Endoscopic carpal tunnel release is a newer procedure that allows division of the transverse carpal ligament with the overlying structures left intact. Use of this procedure purportedly lessens scar formation and allows an earlier return to work and activities of daily living. The wrist is generally splinted for three to four weeks after surgery.

One systematic review³⁴ found equal effectiveness for endoscopic and open surgical techniques in relieving the symptoms of carpal tunnel syndrome, but conflicting evidence on whether endoscopic surgery allows an earlier return to work or other activities. Four studies in the review demonstrated earlier return to activity after endoscopic carpal tunnel release, whereas three studies demonstrated no difference between the techniques.

Complications of surgery include injury to the palmar cutaneous or recurrent motor branch of the median nerve, hypertrophic scarring, laceration of the superficial palmar arch, and tendon adhesion.²⁵ Postoperative infection, hematoma, arterial injury, stiffness, and reflex sympathetic dystrophy are other possible complications.²¹ If carpal tunnel release is incomplete, symptoms may recur.

Incorrect or incomplete diagnosis may be a factor in explaining unsuccessful carpal tunnel release surgery. The strongest predictors of an unfavorable surgical outcome are poor

Both open and endoscopic carpal tunnel release procedures are effective. Whether endoscopic surgery allows an earlier return to work and activities of daily living remains unclear.

scores on patient-reported measures of upper extremity function and mental health status, compensation action connected to carpal tunnel syndrome that involves an attorney, and alcohol use.³²

PREGNANCY

Alterations in fluid balance may predispose some pregnant women to develop carpal tunnel syndrome. Symptoms are typically bilateral and first noted during the third trimester. Conservative measures are appropriate, because symptoms resolve after delivery in most women with pregnancy-related carpal tunnel syndrome.³⁵

The author indicates that he does not have any conflicts of interest. Sources of funding: none reported.

The opinions and assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the U.S. Naval Medical Department or the U.S. Naval Service at large.

REFERENCES

1. Sternbach G. The carpal tunnel syndrome. *J Emerg Med* 1999;17:519-23.
2. Atroshi I, Gummesson C, Johnsson R, Ornstein E, Ranstam J, Rosen I. Prevalence of carpal tunnel syndrome in a general population. *JAMA* 1999;282:153-8.
3. Stevens JC, Witt JC, Smith BE, Weaver AL. The frequency of carpal tunnel syndrome in computer users at a medical facility. *Neurology* 2001;56:1568-70.
4. Stevens JC, Smith BE, Weaver AL, Bosch EP, Deen HG Jr, Wilkens JA. Symptoms of 100 patients with electromyographically verified carpal tunnel syndrome. *Muscle Nerve* 1999;22:1448-56.
5. Von Schroeder HP, Botte MJ. Carpal tunnel syndrome. *Hand Clin* 1996;12:643-55.
6. Stevens JC, Beard CM, O'Fallon WM, Kurland LT. Conditions associated with carpal tunnel syndrome. *Mayo Clin Proc* 1992;67:541-8.

7. D'Arcy CA, McGee S. The rational clinical examination. Does this patient have carpal tunnel syndrome? *JAMA* 2000;283:3110-7.
8. Jablecki CK, Andary MT, Floeter MK, Miller RG, Quartly CA, Vennix MJ, et al. Practice parameter: electrodiagnostic studies in carpal tunnel syndrome. Report of the American Association of Electrodiagnostic Medicine, American Academy of Neurology, and the American Academy of Physical Medicine and Rehabilitation. *Neurology* 2002;58:1589-92.
9. Kanaan N, Sawaya RA. Carpal tunnel syndrome: modern diagnostic and management techniques. *Br J Gen Pract* 2001;51:311-4.
10. Kaplan SJ, Glickel SZ, Eaton RG. Predictive factors in the non-surgical treatment of carpal tunnel syndrome. *J Hand Surg [Br]* 1990;15:106-8.
11. Kruger VL, Kraft GH, Deitz JC, Ameis A, Polissar L. Carpal tunnel syndrome: objective measures and splint use. *Arch Phys Med Rehabil* 1991;72:517-20.
12. Sailer SM. The role of splinting and rehabilitation in the treatment of carpal and cubital tunnel syndromes. *Hand Clin* 1996;12:223-41.
13. Manente G, Torrieri F, Di Blasio F, Staniscia T, Romano F, Uncini A. An innovative hand brace for carpal tunnel syndrome: a randomized controlled trial. *Muscle Nerve* 2001;24:1020-5.
14. Walker WC, Metzler M, Cifu DX, Swartz Z. Neutral wrist splinting in carpal tunnel syndrome: a comparison of night-only versus full-time wear instructions. *Arch Phys Med Rehabil* 2000;81:424-9.
15. Gerritsen AA, de Krom MC, Struijs MA, Scholten RJ, de Vet HC, Bouter LM. Conservative treatment options for carpal tunnel syndrome: a systematic review of randomised controlled trials. *J Neurol* 2002;249:272-80.
16. Spooner GR, Desai HB, Angel JF, Reeder BA, Donat JR. Using pyridoxine to treat carpal tunnel syndrome. Randomized control trial. *Can Fam Physician* 1993;39:2122-7.
17. Chang MH, Chiang HT, Lee SS, Ger LP, Lo YK. Oral drug of choice in carpal tunnel syndrome. *Neurology* 1998;51:390-3.
18. Pal B, Mangion P, Hossain MA, Wallace AS, Diffey BL. Should diuretics be prescribed for idiopathic carpal tunnel syndrome? Results of a controlled trial. *Clin Rehabil* 1988;2:299-301.
19. Stransky M, Rubin A, Lava NS, Lazaro RP. Treatment of carpal tunnel syndrome with vitamin B6: a double-blind study. *South Med J* 1989;82:841-2.
20. Herskovitz S, Berger AR, Lipton RB. Low-dose, short-term oral prednisone in the treatment of carpal tunnel syndrome. *Neurology* 1995;45:1923-5.
21. Whitley JM, McDonnell DE. Carpal tunnel syndrome. A guide to prompt intervention. *Postgrad Med* 1995;97(1):89-92,95-6.
22. Marshall S, Tardif G, Ashworth N. Local corticosteroid injection for carpal tunnel syndrome (Cochrane Review). *Cochrane Database Syst Rev* 2002;(4):CD001554.
23. Wong SM, Hui AC, Tang A, Ho PC, Hung LK, Wong KS, et al. Local vs systemic corticosteroids in the treatment of carpal tunnel syndrome. *Neurology* 2001;56:1565-7.
24. Pfenninger JL. Joint and soft tissue aspiration and injection. In: Pfenninger JL, Fowler GC, eds. *Procedures for primary care physicians*. St. Louis: Mosby, 1994:1036-54.
25. Kulick RG. Carpal tunnel syndrome. *Orthop Clin North Am* 1996;27:345-54.
26. Hunt TR, Osterman AL. Complications of the treatment of carpal tunnel syndrome. *Hand Clin* 1994;10:63-71.
27. Dammers JW, Veering MM, Vermeulen M. Injection with methylprednisolone proximal to the carpal tunnel: randomised double blind trial. *BMJ* 1999;319:884-6.
28. Ebenbichler GR, Resch KL, Nicolakis P, Wiesinger GF, Uhl F, Ghanem AH, et al. Ultrasound treatment for treating the carpal tunnel syndrome: randomised "sham" controlled trial. *BMJ* 1998;316:731-5.
29. Oztas O, Turan B, Bora I, Karakaya MK. Ultrasound therapy effect in carpal tunnel syndrome. *Arch Phys Med Rehabil* 1998;79:1540-4.
30. Longstaff L, Milner RH, O'Sullivan S, Fawcett P. Carpal tunnel syndrome: the correlation between outcome, symptoms and nerve conduction study findings. *J Hand Surg [Br]* 2001;26:475-80.
31. Carter T, Jordan R, Cummins C. Electrodiagnostic techniques in the pre-surgical assessment of patients with carpal tunnel syndrome. A West Midlands Development and Evaluation Service report. Retrieved January 23, 2003, from www.publichealth.bham.ac.uk/wmhtac/pdf/electrodiag.pdf.
32. Katz JN, Losina E, Amick BC III, Fossel AH, Bessette L, Keller RB. Predictors of outcomes of carpal tunnel release. *Arthritis Rheum* 2001;44:1184-93.
33. Gerritsen AA, Uitdehaag BM, van Geldere D, Scholten RJ, de Vet HC, Bouter LM. Systematic review of randomized clinical trials of surgical treatment for carpal tunnel syndrome. *Br J Surg* 2001;88:1285-95.
34. Gerritsen AA, de Vet HC, Scholten RJ, Bertelsmann FW, de Krom MC, Bouter LM. Splinting vs surgery in the treatment of carpal tunnel syndrome: a randomized controlled trial. *JAMA* 2002;288:1245-51.
35. Turgut F, Cetinsahin M, Turgut M, Bolukbasi O. The management of carpal tunnel syndrome in pregnancy. *J Clin Neurosci* 2001;8:332-4.